



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: November 20, 2014

ACCEPTED: December 11, 2014

PUBLISHED: December 29, 2014

# Holographic interpolation between $a$ and $F$

Teruhiko Kawano,<sup>a</sup> Yuki Nakaguchi<sup>a,b</sup> and Tatsuma Nishioka<sup>a</sup>

<sup>a</sup>*Department of Physics, Faculty of Science, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan*

<sup>b</sup>*Kavli Institute for the Physics and Mathematics of the Universe (WPI), The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba 277-8568, Japan*

*E-mail:* [kawano@hep-th.phys.s.u-tokyo.ac.jp](mailto:kawano@hep-th.phys.s.u-tokyo.ac.jp), [yuuki.nakaguchi@ipmu.jp](mailto:yuuki.nakaguchi@ipmu.jp), [nishioka@hep-th.phys.s.u-tokyo.ac.jp](mailto:nishioka@hep-th.phys.s.u-tokyo.ac.jp)

**ABSTRACT:** An interpolating function  $\tilde{F}$  between the  $a$ -anomaly coefficient in even dimensions and the free energy on an odd-dimensional sphere has been proposed recently and is conjectured to monotonically decrease along any renormalization group flow in continuous dimension  $d$ . We examine  $\tilde{F}$  in the large- $N$  CFT's in  $d$  dimensions holographically described by the Einstein-Hilbert gravity in the  $\text{AdS}_{d+1}$  space. We show that  $\tilde{F}$  is a smooth function of  $d$  and correctly interpolates the  $a$  coefficients and the free energies. The monotonicity of  $\tilde{F}$  along an RG flow follows from the analytic continuation of the holographic  $c$ -theorem to continuous  $d$ , which completes the proof of the conjecture.

**KEYWORDS:** AdS-CFT Correspondence, Renormalization Group

**ARXIV EPRINT:** [1410.5973](https://arxiv.org/abs/1410.5973)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Holographic proof of the <math>\tilde{F}</math>-theorem</b>	<b>2</b>

---

## 1 Introduction

A measure of degrees of freedom in a quantum field theory (QFT) remains to be elucidated in arbitrary  $d$  dimensions. Physically, it decreases monotonically as the energy scale is lowered because of the decoupling of massive particles. Implementation of such a measure in any QFT in diverse dimensions is intriguing and desirable to characterize the behavior under a renormalization group (RG) flow.

For even  $d$ , the conformal anomaly in the stress-energy tensor<sup>1</sup>

$$\langle T_{\mu}^{\mu} \rangle = \frac{(-1)^{\frac{d}{2}+1}}{2} a E_d + \sum_i b_i I_i, \quad (1.1)$$

defines the unique  $a$  coefficient for the Euler density  $E_d$  and several  $b_i$  coefficients for the Weyl invariants  $I_i$  labeled by an integer  $i$ . The  $a$  coefficients are believed to be monotonically decreasing along any RG flow, namely the value  $a_{\text{UV}}$  at the ultra-violet (UV) fixed point is equal or greater than that  $a_{\text{IR}}$  at the infra-red (IR) fixed point,  $a_{\text{UV}} \geq a_{\text{IR}}$ . This statement was established in two dimensions by the Zamolodchikov's  $c$ -theorem [1] and in four dimensions by the  $a$ -theorem [2–4]. On the other hand, the  $F$ -theorem asserts that the free energy,  $F \equiv (-1)^{\frac{d-1}{2}} \log Z_{S^d}$ , defined by the conformal invariant partition function  $Z_{S^d}$  on  $S^d$  of radius  $R$ , decreases under any RG flow in odd dimensions [5, 6]. A proof for  $d = 3$  was presented by [7] through the relation of the free energy to the entanglement entropy  $S$  across an entangling surface  $S^{d-2}$  of radius  $R$  in  $\mathbb{R}^{1,d-1}$  [8]

$$F = (-1)^{\frac{d-1}{2}} S, \quad (1.2)$$

that holds for odd  $d$  up to UV divergences.

These two proposals look quite different at first sight, but share the fact that both the  $a$  coefficient and the free energy can be read off on  $S^d$ ; the former arises from the integration of the trace of the stress-energy tensor (1.1) and the latter from the partition function. To interpolate between the  $a$  coefficient and the free energy, Giombi and Klebanov define a new function [9]

$$\tilde{F} \equiv \sin\left(\frac{\pi d}{2}\right) \log Z_{S^d}, \quad (1.3)$$

---

<sup>1</sup>We define the stress-energy tensor by  $T_{\mu\nu} \equiv \frac{2}{\sqrt{g}} \frac{\delta I}{\delta g^{\mu\nu}}$  for an action  $I$ . The Euler density is normalized to be  $\int_{S^d} d^d x \sqrt{g} E_d = 2$ .

which correctly reduces to the free energies for odd  $d$ . They show as  $d$  approaches to even integers<sup>2</sup> (see also [10] as a related work)

$$\tilde{F} = \frac{\pi}{2} a . \quad (1.4)$$

Note that the partition function  $Z_{S^d}$  used in (1.3) is conformal invariant and UV divergent for even  $d$ . The relation (1.4) follows from the fact that the conformal invariant partition function in  $d = 2n + \epsilon$  dimensions behaves as  $\log Z_{S^d} = (-1)^{\frac{d}{2}} \frac{a}{2\epsilon} + O(1)$  for small  $\epsilon$ . This is because one has to add a local counter term

$$I_{\text{c.t.}} = (-1)^{\frac{d}{2}+1} \frac{a}{2\epsilon} \int_{S^d} d^d x \sqrt{g} E_{2n} , \quad (1.5)$$

to the partition function to obtain the renormalized partition function  $\log Z_{S^d}^{(\text{ren})} = \log Z_{S^d} + I_{\text{c.t.}}$ , reproducing the conformal anomaly  $\log Z_{S^{2n}}^{(\text{ren})} = (-1)^{n+1} a \log R$  on  $S^{2n}$  of radius  $R$  in  $\epsilon \rightarrow 0$  limit.

The function  $\tilde{F}$  is also defined for non-integer  $d$  and therefore smoothly interpolates between the  $a$  coefficients in even dimensions and the free energies in odd dimensions. They conjecture that  $\tilde{F}$  is positive and decreases along any RG flow in arbitrary  $d$  dimensions, based on several examples including a double-trace deformation of the large- $N$  conformal field theory (CFT). We will call their proposal the  $\tilde{F}$ -theorem.

In this letter, we provide a further evidence to the  $\tilde{F}$ -theorem from the holographic viewpoint. To this end, we take advantage of the relation (1.2) and calculate the holographic entanglement entropy [11, 12] across a sphere  $S^{d-2}$  in the Einstein-Hilbert gravity on the  $\text{AdS}_{d+1}$  space. We perform the dimensional regularization in the bulk and obtain the analytic result of  $\tilde{F}$  that is a positive and smooth function of dimension  $d$ . We show that the equality (1.4) holds for even  $d$  and furthermore prove the  $\tilde{F}$ -theorem that follows from the holographic  $c$ -theorem [13–16] assuming the dimensional continuation of the null energy condition.

## 2 Holographic proof of the $\tilde{F}$ -theorem

We will evaluate  $\tilde{F}$  with the relation (1.2) between the free energy on  $S^d$  and the entanglement entropy across  $S^{d-2}$ . The latter can be holographically calculated by the Ryu-Takayanagi formula in the Einstein-Hilbert gravity [11, 12]

$$S = \frac{\text{Area}(\gamma)}{4G_N^{(d+1)}} , \quad (2.1)$$

where  $G_N^{(d+1)}$  is the Newton constant, and  $\gamma$  stands for the  $(d-1)$ -dimensional minimal surface in the  $\text{AdS}_{d+1}$  space, whose boundary is the entangling surface  $S^{d-2}$ . Since the boundary of the  $\text{AdS}_{d+1}$  space is the flat space  $\mathbb{R}^{1,d-1}$ , we will use the Poincaré coordinates

$$ds^2 = L^2 \frac{dz^2 - dt^2 + dr^2 + r^2 d\Omega_{d-2}^2}{z^2} , \quad (2.2)$$

---

<sup>2</sup>There is no sign factor  $(-1)^{d/2}$  in the right hand side because our convention of the  $a$ -anomaly (1.1) differs from theirs in [9].

where  $L$  is the AdS radius. The entangling surface is located at  $t = 0$  and  $r = R$  at the boundary  $z = 0$ . In these coordinates, the minimal surface  $\gamma$  in the bulk is a hemi-hypersphere satisfying  $r^2 + z^2 = R^2$  [11, 12]. This solution leads the entanglement entropy across  $S^{d-2}$

$$S = \frac{1}{4G_N^{(d+1)}} L^{d-1} \text{Vol}(S^{d-2}) \int_{\epsilon/R}^1 dy \frac{(1-y^2)^{\frac{d-3}{2}}}{y^{d-1}}, \quad (2.3)$$

where we introduced a small cutoff at  $z = \epsilon$  to regularize the UV divergence and  $\text{Vol}(S^{d-2})$  is the volume of a unit  $(d-2)$ -dimensional round sphere. Expanding the integrand with respect to  $y$  and performing the integration, one obtains the UV divergent parts of the entanglement entropy. We, however, want to employ the dimensional regularization instead of putting the UV cutoff at  $z = \epsilon$  for our purpose. So we take  $\epsilon = 0$  and carry out the integral in the range  $1 < d < 2$ , that yields

$$S = \frac{L^{d-1}}{4G_N^{(d+1)}} \pi^{\frac{d}{2}-1} \Gamma\left(1 - \frac{d}{2}\right). \quad (2.4)$$

Then we analytically continue  $d$  to any real value. It is clear that there are poles at even  $d$  in the entanglement entropy (2.4) corresponding to the conformal anomalies. Finally, using the relations (1.2) and (1.3), and the formula  $\Gamma(z)\Gamma(1-z) = \pi/\sin(\pi z)$ , we obtain  $\tilde{F}$  in the holographic theories

$$\tilde{F} = \frac{L^{d-1}}{4G_N^{(d+1)}} \frac{\pi^{\frac{d}{2}}}{\Gamma\left(\frac{d}{2}\right)}. \quad (2.5)$$

This is manifestly a positive and smooth function of dimension  $d$  without poles at even  $d$ .

Now let us extrapolate the holographic values of  $\tilde{F}$  to even dimensions and see if the relation (1.4) holds. The  $a$  coefficients holographically computed in the Einstein-Hilbert gravity are known to be [15–18]

$$a = \frac{L^{d-1}}{2\pi G_N^{(d+1)}} \frac{\pi^{\frac{d}{2}}}{\Gamma\left(\frac{d}{2}\right)}. \quad (2.6)$$

Combining it with (2.5), we confirm the relation (1.4) between  $\tilde{F}$  and  $a$ . Moreover, imposing the null energy condition in the bulk, the holographic  $c$ -theorem states that the  $a$  coefficient given by (2.6) satisfies the monotonicity,  $a_{\text{UV}} \geq a_{\text{IR}}$ , for positive integer  $d$  [13–16]. Assuming the analytic continuation of dimension  $d$  in the gravity, the holographic  $c$ -theorem holds for  $d \geq 1$ ,<sup>3</sup> which assures the  $\tilde{F}$ -theorem due to the relation (1.4).

## Acknowledgments

We are grateful to Y. Tachikawa and K. Yonekura for valuable discussions and to S. Giombi and I. Klebanov for correspondence. The work of T. K. was supported in part by a Grant-in-Aid #23540286 from the MEXT of Japan. The work of Y. N. was supported in part by JSPS Research Fellowship for Young Scientists and World Premier International Research Center Initiative (WPI) from the MEXT of Japan.

<sup>3</sup>The null energy condition  $T_{\mu\nu}\xi^\mu\xi^\nu \geq 0$  is crucial in the proof of the holographic  $c$ -theorem [14–16] where the  $d$ -dimensional null vector  $\xi$  has only two non-zero components  $\xi^z$  and  $\xi^t$ . Thus defining a formal null vector  $\xi = (\xi^z, \xi^t, 0, \dots, 0)$  in continuous  $d$  dimensions, the proof can be carried over for  $d \geq 1$ .

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] A.B. Zamolodchikov, “Irreversibility” of the flux of the renormalization group in a 2D field theory, *JETP Lett.* **43** (1986) 730 [*Pisma Zh. Eksp. Teor. Fiz.* **43** (1986) 565] [[INSPIRE](#)].
- [2] J.L. Cardy, Is there a  $c$ -theorem in four dimensions?, *Phys. Lett. B* **215** (1988) 749 [[INSPIRE](#)].
- [3] Z. Komargodski and A. Schwimmer, On Renormalization Group Flows in Four Dimensions, *JHEP* **12** (2011) 099 [[arXiv:1107.3987](#)] [[INSPIRE](#)].
- [4] Z. Komargodski, The Constraints of Conformal Symmetry on RG Flows, *JHEP* **07** (2012) 069 [[arXiv:1112.4538](#)] [[INSPIRE](#)].
- [5] D.L. Jafferis, I.R. Klebanov, S.S. Pufu and B.R. Safdi, Towards the  $F$ -Theorem:  $N = 2$  Field Theories on the Three-Sphere, *JHEP* **06** (2011) 102 [[arXiv:1103.1181](#)] [[INSPIRE](#)].
- [6] I.R. Klebanov, S.S. Pufu and B.R. Safdi,  $F$ -Theorem without Supersymmetry, *JHEP* **10** (2011) 038 [[arXiv:1105.4598](#)] [[INSPIRE](#)].
- [7] H. Casini and M. Huerta, On the RG running of the entanglement entropy of a circle, *Phys. Rev. D* **85** (2012) 125016 [[arXiv:1202.5650](#)] [[INSPIRE](#)].
- [8] H. Casini, M. Huerta and R.C. Myers, Towards a derivation of holographic entanglement entropy, *JHEP* **05** (2011) 036 [[arXiv:1102.0440](#)] [[INSPIRE](#)].
- [9] S. Giombi and I.R. Klebanov, Interpolating between  $a$  and  $F$ , [arXiv:1409.1937](#) [[INSPIRE](#)].
- [10] K. Yonekura, Perturbative  $c$ -theorem in  $d$ -dimensions, *JHEP* **04** (2013) 011 [[arXiv:1212.3028](#)] [[INSPIRE](#)].
- [11] S. Ryu and T. Takayanagi, Holographic derivation of entanglement entropy from AdS/CFT, *Phys. Rev. Lett.* **96** (2006) 181602 [[hep-th/0603001](#)] [[INSPIRE](#)].
- [12] S. Ryu and T. Takayanagi, Aspects of Holographic Entanglement Entropy, *JHEP* **08** (2006) 045 [[hep-th/0605073](#)] [[INSPIRE](#)].
- [13] L. Girardello, M. Petrini, M. Porrati and A. Zaffaroni, Novel local CFT and exact results on perturbations of  $N = 4$  super Yang-Mills from AdS dynamics, *JHEP* **12** (1998) 022 [[hep-th/9810126](#)] [[INSPIRE](#)].
- [14] D.Z. Freedman, S.S. Gubser, K. Pilch and N.P. Warner, Renormalization group flows from holography supersymmetry and a  $c$  theorem, *Adv. Theor. Math. Phys.* **3** (1999) 363 [[hep-th/9904017](#)] [[INSPIRE](#)].
- [15] R.C. Myers and A. Sinha, Seeing a  $c$ -theorem with holography, *Phys. Rev. D* **82** (2010) 046006 [[arXiv:1006.1263](#)] [[INSPIRE](#)].
- [16] R.C. Myers and A. Sinha, Holographic  $c$ -theorems in arbitrary dimensions, *JHEP* **01** (2011) 125 [[arXiv:1011.5819](#)] [[INSPIRE](#)].
- [17] C. Imbimbo, A. Schwimmer, S. Theisen and S. Yankielowicz, Diffeomorphisms and holographic anomalies, *Class. Quant. Grav.* **17** (2000) 1129 [[hep-th/9910267](#)] [[INSPIRE](#)].
- [18] A. Schwimmer and S. Theisen, Entanglement Entropy, Trace Anomalies and Holography, *Nucl. Phys. B* **801** (2008) 1 [[arXiv:0802.1017](#)] [[INSPIRE](#)].